

УДК 632.4

**A BIOINFORMATICS CHARACTERIZATION OF THE RNA-DEPENDENT RNA POLYMERASE FROM THE PARASITIC FUNGI, *Metarhizium anisopliae***

**C. N. WEIR<sup>1,2</sup>**

**M. Mol. Biol. and Ph. D candidate**

**E. WONG<sup>3</sup> Ph. D.**

**G. F. KING<sup>4</sup> Ph. D.**

<sup>1</sup> *Walter and Eliza Hall Institute of Medical Research, 1G Royal Parade, Parkville 3052, Victoria, Australia, e-mail: [weir.c@wehi.edu.au](mailto:weir.c@wehi.edu.au)*

<sup>2</sup> *University of Melbourne, Department of Medical Biology, Parkville 3010, Victoria, Australia*

<sup>3</sup> *European Bioinformatics Institute (EMBL-EBI), Wellcome Trust Genome Campus, Hinxton, Cambridgeshire CB10 1SD, United Kingdom*

<sup>4</sup> *Institute for Molecular Bioscience, University of Queensland, St. Lucia 4072, Brisbane, Queensland, Australia*

***Metarhizium anisopliae* is an entomopathogenic organism that is parasitic towards insects, including those that serve as vectors for human disease, including malaria parasite infected *Anopheles* mosquitoes. Directed bio-engineering of parasitic *M. anisopliae* has been proposed as a potentially novel vector control strategy. However, many parasitic fungi possess a RNA-dependent RNA polymerase (*RdRp*) gene that serves as a defence against transgenes. Here we characterize the *RdRp* gene of *M. anisopliae*, including the context of the gene in relation to surrounding genes and theoretical modeling based on the *RdRp* enzyme of *Neurospora crassa*.**

Keywords: entomopathogenic fungi, *Metarhizium anisopliae*, parasites, transgenes, vector control.

In fungi, including parasitic *Cordyceps* species and some of the *Metarhizium* species, RNA-dependent RNA polymerases (RdRp) are involved in the phenomenon of gene-silencing, or quelling, where they act at the post-transcriptional level, affecting both endogenous genes and transgenes<sup>1</sup>. Slight differences in molecular pathways exist depending on what type of gene is involved, however here we will focus on exogenous transgenes (for a review of both mechanisms, please refer to Dang Y et al.<sup>2</sup>). During quelling, a diffusible trans-acting aberrant RNA (abRNA) molecule of the expressed gene is produced and converted to double-stranded RNA (dsRNA) via RdRp's activity allowing for the synthesis of complementary RNA molecules<sup>1-3</sup>. The dsRNA is recognized by the Dicer enzyme (or Dicer-like proteins), degrading the dsRNA into small-interfering RNA (siRNA) molecules that get incorporated into the RNA-induced silencing complex (RISC)<sup>1,2</sup>. RISC uses the siRNA's to guide nuclease activity



### **Protein Blast and Domain Structure**

Protein blast searches were performed using the NCBI BLASTp function; BLAST searches were first performed only against fungi then against solved PDB entries (with the latter generating the information regarding *Neurospora crassa*). Domain architecture information was generated with both NCBI (<http://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi?and>) and Pfam (<http://pfam.sanger.ac.uk/protein/E9EKW8>). The sequence alignment between *N. crassa* and *M. anisopliae* RdRp was manually edited in order to improve the alignment. This was checked with the associated PDB file for PDB accession 2J7N. Pymol software used for modeling.

### **Homology**

Fungi analyzed were selected based upon BLASTp results and those related in the literature<sup>9</sup>. For protein selection, the highest E-value results were used and hypothetical and predicted proteins were excluded unless, upon analyzing their information on NCBI, they showed evidence for being a true RdRp, such as similar sequences. Multiple sequence alignments were performed using Muscle (<http://www.ebi.ac.uk/Tools/msa/muscle/>). This information was used with MEGA 5.05 software including Muscle alignments, and phylogenetic tools. A phylogenetic neighbour joining tree was generated with the Bootstrap method (500 bootstraps used), Poisson modeling and pair wise deletion for the data treatment and amino acid substitutions allowed in the model. It should be noted that whilst *M. acridium* is highly related to *M. anisopliae*, it was excluded from the phylogenetic analysis of RdRp since the sequence that currently exists for it is incomplete (accession EFY87605).

## **Results and discussion**

### **Gene Structure**

The genome and transcriptome of *M. anisopliae* had been sequenced by Gao Q et al.<sup>9</sup> and genes curated using EMBL-EBI software. The genome was shotgun-sequenced into 1271 contigs with each loci tagged as MAA; the *RdRp* gene (MAA\_00977) is located on contig ADNJO1000234.1 (c234)<sup>9</sup>. After searching EMBL-EBI for «*M. anisopliae* ARSEF 23 RNA-dependant RNA polymerase» relevant gene information was obtained. Sequence length was reported as 3345 base pairs (minus the 222bp of 3 introns), comprised of 4 exons corresponding to the predicted 1114 amino acids. The transcriptional start site is coded for by amethionine, and the stop codon sequence is TAG. No promoters were described, so an attempt was made to find possible ones using Transfac (see Methods). A FASTA formatted sequence 200 bp upstream (towards the 5' region) of *RdRp*'s start codon was input to the Transfac's Patch 1.0 tool, which uses an algorithm for pattern-based prediction of transcription factor binding sites. A likely promoter site, GAGTCA, was found to be 31 bp upstream of the aforementioned start codon. With a score of 100 and no mismatches, this corresponded to a region where the binding factor characterized in *Saccharomyces cerevisiae*, GCN4, interacts.

### **Gene Context**

Currently, the chromosome location of *RdRp* is unknown; only the contig location c234 exists. Gene prediction software (Fgenesh by Softberry) was used to predict adjacent genes along c234. EMBL-EBI software gave information

pertaining to locus tags (identifiers which are applied systematically to all genes in a genome within the context of sequencing projects) for genes within each contig. Using this, the overall context of *RdRp* and its surrounding characterized genes could be viewed, along with contig assembly. The two closest downstream genes from *RdRp*'s stop codon, reported from the gene organization within the contigs, are those coding for the C6 transcription factor *GliZ2* and a protein efflux pump, which are located 6477 and 12638 nucleotides away respectively. The nearest upstream gene, which is 9885 nucleotides away from the start codon of *RdRp*, codes for a hypothetical protein.

Given the distance from *RdRp*, both for upstream and downstream genes, a manual search for closer genes was carried out. As Figure 1 reveals, two such genes were identified either side of *RdRp*; a peroxidase protein 1943 nucleotides upstream of *RdRp*'s start codon, and the NFX1-type zinc finger-containing protein 1, which is 569 nucleotides downstream of *RdRp*'s stop codon. This figure, approximately to scale, shows the improved gene context of *RdRp* with respect to its surrounding genes after manually searching. Directionality of transcription is shown as arrows above the genes, with their size (including introns) and nucleotide location on the contig given below.




**Figure 1.** Gene context of *RdRp*:

PP – peroxidase protein; NFX1 – NFX1-type zinc finger-containing protein. All gene sizes shown are inclusive of introns and exons

### ***Protein Blast and Domain Structure***

A search of the conserved domain database at NCBI and the Pfam database using the amino acid sequence of *RdRp*, was performed (see Methods). The results are shown in Figure 2A and 2B. Both NCBI and Pfam revealed a single characterized domain for *RdRp*, corresponding to amino acid residues 314–925. The Pfam database indicates this family of proteins are important for eukaryotic post-transcriptional gene silencing and that there is a core catalytic domain responsible for the activity. Comparisons of the sequence similarity between this domain and the *RdRp* domain in the orthologous protein from the fungus *Neurospora crassa*. *N. crassa* was used for two reasons; firstly, the *RdRp* of *N. crassa* has been well characterized including a complete crystal structure<sup>10</sup>. Secondly, as Gao Q et al. shows<sup>9</sup>, they are both related at the class level. There is 30 % identity between a 333-residue stretch within the given domain and the active catalytic sub-domain from the *N. crassa* protein (Figure 2C).

Despite low overall sequence homology, the catalytic aspartate residues are conserved between the two species<sup>10,11</sup>. The catalytic aspartate residues interact with a magnesium cation and are responsible for the selection of ribonucleoside triphosphates incorporated into the incoming abRNA, making it dsRNA<sup>10,12</sup>. These conserved aspartate residues are shown in Figure 2C at residues 1007, 1009 and 1011 in the sequence of *N. crassa* *RdRp*.




**Figure 2.** BLASTp and domain information:

**A** – The top two hits after blasting *M. anisopliae* RdRp against known fungal proteins. RdRp proteins from *Cordyceps militaris* and *Verticillium dahlia* showed highest sequence coverage and maximum sequence identity to *M. anisopliae* RdRp; **B** – Domain architecture of *M. anisopliae* RdRp showing the 611-residue domain, compared to the catalytic sub-domain of *N. crassa* RdRp. The figure was made using both NCBI and Pfam data; **C** – Sequence alignment of residues from (B) showing that *M. anisopliae* RdRp has 30 % sequence identity over 333 residues to *N. crassa* RdRp with an E-value of 1.34e-18. Residues in orange are homologous

## Homology

The RdRp gene is found in viruses, plants, protozoa, fungi, and nematodes, although it is absent in insects or vertebrates. There are differences among these organisms with respect to domain organization which results in a catalytic domain possessing either right handed or double-barrelled structural organization<sup>10</sup>. Fungi, in contrast to viruses, possess the latter organization<sup>10</sup>, which will be discussed below in the Additional Information section. Figure 3 shows the phylogenetic relationship between various fungal species and their RdRp protein. The organization of fungi shown in Figure 3 corresponds to the published phylogenetic tree (see Gao Q et al.<sup>9</sup>), showing inferred evolutionary relationships between each fungus.




**Figure 3.** Phylogenetic relationships between fungal RdRps

This neighbour-joining tree, made using MEGA 5.05 (see Methods), demonstrates the evolutionary relationships between nine fungal RdRp proteins based on amino acid sequences. There are three main clades, which match overall the organization seen amongst genomes of the same fungi<sup>9</sup>. Clade 1 contains *M. anisopliae*, *C. militaris*, *V. dahlia*, *N. crassa*, *A. nidulans* and *M. oryzae*. Clade 2 contains *B. fuckeliana* and *S. sclerotiorum*. Clade 3 contains *S. pombe*. Taxonomic grouping is also demonstrated. The cut-off for significance regarding bootstrap values was anything below 65. The Poisson model was used in MEGA 5.05's phylogeny tool and 500 bootstraps were selected.

## Visualization of the Catalytic Region of RdRp

No 3D structure currently exists for *M. anisopliae* RdRp, however there is one for *N. crassa* RdRp, as shown on the PDB under the identifying number 2J7N. It was decided, given the reasonable level of sequence identity between the two sequences shown in Figure 2C, that identification of the residues in the catalytic region of *M. anisopliae* RdRp should be carried out (Figure 4). This revealed the catalytic site with the predicted «catalytic» arginine residues shown interacting with a magnesium cation. These results show that all of the key residues reported in the literature with respect to the catalytic activity of *N. crassa* RdRp are conserved in the RdRp of *M. anisopliae*. The tyrosine residue at position 1010, which is sandwiched between the catalytic residues D1009 and D1011 in *N. crassa* RdRp (Figure 4), is substituted for a leucine in *M. anisopliae* RdRp. Given that both of these residues possess relatively large hydrophobic side-chains, it is unlikely that this substitution would have any negative effects on catalytic activity.





**Figure 4.** Catalytic site of RdRp

The crystallographic model solved to 2.3 Å of *N. crassa* RdRp was used as the model of homology (PDB under accession number 2J7N). Key residues that are highly conserved among all known RdRps are labeled, and those that are identical in *M. anisopliae* RdRp are colored cyan (the blue are found in *N. crassa* RdRp). The three homologous catalytic aspartic acid residues are labeled and colored pink, and are shown interacting with the magnesium cation that lies between each DPBB. There is a tyrosine residue at amino acid position 1010 (not labeled, but is shown in blue) which in the corresponding sequence in *M. anisopliae* is a leucine. The distance of the interaction between each catalytic aspartic acid residue and the magnesium cation is between 2.1 and 2.2 Å.

RdRp enzymes are found in a variety of both eukaryotic and prokaryotic organisms and serve as regulators of gene expression at the post-transcriptional level<sup>10-12</sup>. Given our future research directions involving transgene incorporation into *M. anisopliae* to improve its efficacy for controlling insect vectors of human disease, and the fact that transgenes have been known to initiate gene silencing, it is important to characterize the mechanisms behind this<sup>2,5,6</sup>. This report has elaborated on the gene silencing protein, RdRp, in *M. anisopliae* via the use of bioinformatic tools, with several key findings made.

During *RdRp* gene characterization, the finding that the promoter site GAGTCA is the sequence for binding factor GCN4 was of interest. GCN4 has been associated with the activation of genes involved in protein and purine synthesis, and expressed during times of stress<sup>14</sup>. One could consider the introduction and expression of transgenes to be a stressor to cells since indeed this has been reported in plants with similar silencing mechanisms to fungi, which evolved to possess viral defence mechanisms<sup>1,15</sup>. It therefore makes sense that this region would lie near *RdRp*.

Despite the fairly low overall sequence identity of 30 % between *M. anisopliae* and *N. crassa* RdRp, the conservation of key residues in the catalytic site was expected as predicted from the literature<sup>10,11</sup>. Modeling and matching identical residues was a good way of showing this and relating the form to function (Figure 4). The sequences lying outside of the domain given in Figure 3B for *M.*

*anisopliae* RdRp may code for areas of the protein such as the slab, head or neck regions of the full protein.

The identification and analysis of the *M. anisopliae* RdRp gene reveals that there is likely to be a fully functioning molecular mechanism involved in gene silencing in *M. anisopliae*. This paves the way for development of knockout strains (such as the *qde-1* knockout of *N. crassa*) that may have improved transgene expression<sup>3</sup>. The time taken (20 days) for the transgenic *Metarhizium* engineered by Fang et al. to kill *Plasmodium*-infected mosquitoes may have been reduced with the use of a *RdRp* knockout, if indeed transgene silencing was occurring<sup>6</sup>.

In the future, fluorescent *in situ* hybridization could be used to locate the chromosome on which *RdRp* is found<sup>16</sup>. Additionally, it would be of interest to work out the crystal structure of *M. anisopliae* RdRp since it is gaining popularity for use as an entomopathogenic agent.

### Literature

1. Nakayashiki H. RNA silencing in fungi: Mechanisms and applications // FEBS Letters. – 2005. – V. 579. – P. 5950–5957.
2. Dang Y. et al. RNA Interference in fungi: Pathways, functions and applications // Eukary Cell. – 2011. – V. 10. – P. 1148–1155.
3. Cogoni C., Macino G. Gene silencing in *Neurospora crassa* requires a protein homologous to RNA-dependant RNA polymerase // Nature. – 1999. – V. 399. – P. 166–169.
4. Cogoni C. Homology-dependant gene silencing mechanisms in fungi // Annu. Rev. Microbiol. – 2001. – V. 55. – P. 381–406.
5. Fang W., Azimzadeh P., St. Leger R.J. Strain improvement of fungal insecticides for controlling insect pests and vector-borne diseases // Curr. Opin. Microbiol. – 2012. – V. 15. – P. 1–7.
6. Fang W. et al. Development of transgenic fungi that kill human malaria parasites in mosquitoes // Science. – 2011. – V. 331. – P. 1074–1077.
7. Wang C., St. Leger R.J. A scorpion neurotoxin increases the potency of a fungal insecticide // Nature Biotech. – 2007. – V. 25. – P. 1455–1456.
8. Saez N.J. et al. Spider-venom peptides as therapeutics // Toxins. – 2010. – V. 2. – P. 2851–2871.
9. Gao Q. et al. Genome sequencing and comparative transcriptomics of the model entomopathogenic fungi *Metarhizium anisopliae* and *M. acridium* // PLoS Genetics. – 2011. – V. 7. – P. 100–126.
10. Salgado P.S. et al. The structure of an RNAi polymerase links RNA silencing and transcription // PLoS Biol. – 2006. – V. 4. – e434.
11. Gohara D.W. et al. Poliovirus RNA-dependent RNA polymerase (3Dpol): structural, biochemical, and biological analysis of conserved structural motifs A and B // J. Biol. Chem. – 2000. – V. 275. – P. 25523–25532.
12. O'Reilly E.K., Kao C.C. Analysis of RNA-dependent RNA polymerase structure and function as guided by known polymerase structures and computer predictions of secondary structure // Virology. – 1998. – V. 252. – P. 287–303.
13. Kupfer D.M. et al. Introns and splicing elements of five diverse fungi // Eukary Cell. – 2004. – V. 3. – P. 1088–10100.
14. Mascarenhas C. et al. Gcn4 is required for the response to peroxide stress in the yeast *Saccharomyces cerevisiae* // MBoC. – 2008. – V. 19. – P. 2995–3007.
15. Vance V., Vaucheret H. RNA silencing in plants – defense and counter defense // Science. – 2001. – V. 292. – P. 2277–2280.



16. Cremer T., Cremer C. Chromosome territories, nuclear architecture and gene regulation in mammalian cells // Nature Rev Genetics. – 2001. – V. 2. – P. 292–302.

**Биоинформационная оценка РНК-зависимой РНК-полимеразы  
гриба-паразита *Metarhizium anisopliae***

**К. Н. Вейер<sup>1,2</sup>, Э. Ронг<sup>3</sup>, Г. Ф. Кинг<sup>4</sup>**  
**кандидаты биологических наук**

<sup>1</sup> Институт Медицинских Исследований Уолтера и Элизы Холл, 1G Ройал  
Парад, Парквилл 3052, Виктория, Австралия

<sup>2</sup> Мельбурнский университет, Отдел Медицинской Биологии, Парквилл  
3010, Виктория, Австралия

<sup>3</sup> Европейский институт биоинформатики (EMBL-EBI), Кампус "Wellcome  
Trust Genome", Хинкстон, Кембриджшир CB10 1SD, Великобритания

<sup>4</sup> Институт молекулярной биологии, Университет Квинсленда, Св. Лючия  
4072, Брисбейн, Квинслэнд, Австралия

*Metarhizium anisopliae* представляет собой энтомопатогенный гриб, паразитирующий на насекомых, включая тех, которые являются переносчиками инвазии, в том числе, зараженных малярийными паразитами москитов *Anopheles*. В качестве потенциально новой стратегии контроля переносчиков при исследовании *M. anisopliae* предлагается использовать биоинженерный метод. Однако, многие грибы-паразиты содержат ген РНК-зависимой РНК-полимеразы (RdRp), который служит защитой от трансгенов. Дана характеристика гена *RdRp* гриба *M. anisopliae*, включая его строение, представленное во взаимосвязи с окружающими его генами, и генетическое моделирование, основанное на изучении RdRp белка гриба *Neurospora crassa*.

Ключевые слова: энтомопатогенные грибы, *Metarhizium anisopliae*, паразиты, трансгены, контроль переносчика.